

A List of $1 + 1$ Dimensional Integrable Equations and Their Properties

Jing Ping WANG

Division of Mathematics & Computer Science, Faculty of Sciences, Vrije Universiteit, Amsterdam, The Netherlands

E-mail: wang@cs.vu.nl

Received August 29, 2001; Revised October 26, 2001; Accepted 1 November, 2001

Abstract

This paper contains a list of known integrable systems. It gives their recursion-, Hamiltonian-, symplectic- and cosymplectic operator, roots of their symmetries and their scaling symmetry.

1 Introduction

In this paper we give a list of known integrable systems. This list is far from being complete, even if we restrict ourselves to

- What is known today,
- Systems with less than four components, and
- Systems referred to in the literature with a name.

Originally this list was part of the author's thesis [Wan98]. Since this thesis only has a limited distribution and the reactions to this list were very favorable, we decided to publish it in a more accessible way.

The theoretical background for our list can be found in [Olv93, Dor93]. A more encyclopedic approach to this subject is taken in [Ibr96], where lists of integrable equations with their properties, mainly special solutions, Lie symmetries and conservation laws, are given. For every equation we aim to give a table containing:

- the equation itself,
- its cosymplectic operator (Hamiltonian operator, cf. [Mag78]),
- its Hamiltonian function corresponding to the cosymplectic operator,
- its symplectic operator,
- its recursion operator (possibly resulting from the cosymplectic and symplectic operators, cf. [FF80, FF81]), or its master symmetry,

Copyright © 2002 by J P Wang

- roots of its symmetries,
- its scaling symmetry.

There are also other important properties such as Miura transformations between the systems and Lax pair (cf. [Lax68]), but these are mainly ignored here. We refer to the source where we learned about each system; no attempt has been made to track the historical origins. At this point, we do not claim any new results though we did put in an effort to compute all the subjects in the list; the decomposition of the recursion operator in symplectic and cosymplectic operators mainly relies on the theory of recursion operators (cf. [SW01a]).

We hope that this material can serve as a source of motivation for future research, since it allows one to quickly formulate or dismiss general statements.

First we give the definitions of the subjects in our lists.

Definition 1. *Given an evolution equation $u_t = K(x, t, u, \dots, u_n)$, where $u_i = \frac{\partial^i u}{\partial x^i}$, we define*

$$\begin{aligned} h \text{ is a symmetry if } & \mathcal{L}_K h = \frac{\partial h}{\partial t} + D_h[K] - D_K[h] = 0, \\ \mathfrak{H} \text{ is a cosymplectic operator if } & \mathcal{L}_K \mathfrak{H} = \frac{\partial \mathfrak{H}}{\partial t} + D_{\mathfrak{H}}[K] - D_K \mathfrak{H} - \mathfrak{H} D_K^* = 0, \\ \mathfrak{J} \text{ is a symplectic operator if } & \mathcal{L}_K \mathfrak{J} = \frac{\partial \mathfrak{J}}{\partial t} + D_{\mathfrak{J}}[K] + \mathfrak{J} D_K + D_K^* \mathfrak{J} = 0, \\ \mathfrak{R} \text{ is a recursion operator if } & \mathcal{L}_K \mathfrak{R} = \frac{\partial \mathfrak{R}}{\partial t} + D_{\mathfrak{R}}[K] - D_K \mathfrak{R} + \mathfrak{R} D_K = 0, \end{aligned}$$

where \mathcal{L}_K denotes the Lie derivative and \star means conjugation. Moreover, for the operators, the formulae are only valid on the domain of the left hand sides of the identities. A symmetry is called a **root** of a hierarchy if it is in the domain, but not in the image, of the recursion operator. A vectorfield $\Sigma = f(t)\partial_t + S\partial_u$ is called a **scaling symmetry** of the equation if $\mathcal{L}_{\Sigma}K = \lambda K, \lambda \in \mathbb{C}$.

Remark 2. *Here we still use the standard definition of recursion operator in the literature. We refer the reader to [SW01b], where the term weak recursion operator is coined, for a discussion of the problems with this definition.*

Acknowledgement. *The author would like to thank Dr. Jan Sanders, Vrije Universiteit Amsterdam, for his strong encouragement to carry out this work and Prof. A.G. Meshkov, Oryol State University, Russia, for correcting several typographical errors.*

2 List of integrable systems and some of their properties

2.1 Burgers' equation

Reference: [Olv93, p. 315], [Oev84, p. 38];

Equation	$u_t = u_2 + uu_1$	
Hamiltonian	None	[Fuc79]
Cosymplectic	None	[FF81]
Symplectic	None	[FF81]
Recursion	$\mathfrak{R}_1 = D_x + \frac{1}{2}u + \frac{1}{2}u_1 D_x^{-1}$ $\mathfrak{R}_2 = tD_x + \frac{1}{2}(tu + x) + \frac{1}{2}(tu_1 + 1)D_x^{-1}$	[Olv77] [Olv93]
Root	$u_1, tu_1 + 1$	\star
Scaling	$-2t\partial_t + (xu_1 + u)\partial_u$	

★ Since \mathfrak{R}_1 and \mathfrak{R}_2 are both recursion operators of the equation, we obtain a double infinity of the symmetries, by applying \mathfrak{R}_1 or \mathfrak{R}_2 successively to u_1 and $tu_1 + 1$. Note that since $\mathfrak{R}_1\mathfrak{R}_2 = \mathfrak{R}_2\mathfrak{R}_1 + \frac{1}{2}$ and $\mathfrak{R}_1(tu_1 + 1) = \mathfrak{R}_2(u_1)$, if we are only interested in independent symmetries, it does not matter in which order \mathfrak{R}_1 and \mathfrak{R}_2 are applied.

Notice that there is a difference between the root of an operator and the root of symmetries for an equation. For \mathfrak{R}_1 , we can take $tu_1 + 1$ as a root of symmetries for Burgers' equation since $\mathfrak{R}_1 = D_x + \frac{1}{2}D_x(uD_x^{-1}\cdot)$, but it is not a root of \mathfrak{R}_1 since it is not its symmetry.

We refer to [Ma93] for coupled Burgers systems.

2.2 Potential Burgers' equation

Reference: [Olv93, pp. 311, 317];

Equation	$u_t = u_2 + u_1^2$	
Hamiltonian	None	[Fuc79]
Cosymplectic	None	[FF81]
Symplectic	None	[FF81]
Recursion	$\mathfrak{R}_1 = D_x + u_1$ $\mathfrak{R}_2 = tD_x + tu_1 + \frac{1}{2}x$	[Olv93]
Root	1	
Scaling	xu_1	[Olv93]

The same arguments hold here as Burgers' equation since $\mathfrak{R}_1\mathfrak{R}_2 = \mathfrak{R}_2\mathfrak{R}_1 + \frac{1}{2}$.

The author of [Fok80] found the 2nd-order equations of the form $u_t = u_2 + f(u, u_1)$, which possess a 3rd-order symmetry and obtained the following equations:

$$u_t = u_2 + \frac{f''(u)}{f'(u)}u_1^2 + \alpha f(u)u_1, \quad (2.1)$$

where α is constant and $f(u)$ is an arbitrary function, with the recursion operator $D_x + \frac{f''(u)}{f'(u)}u_1 + \frac{1}{2}\alpha f(u) + \frac{1}{2}\alpha u_1 D_x^{-1}f'(u)$.

$$u_t = u_2 + \frac{\gamma - f'(u)}{f(u)}u_1^2 + \alpha f(u), \quad (2.2)$$

where $f(u)$ is an arbitrary function and α, γ are constant, with the recursion operator $D_x + \frac{\gamma - f'(u)}{f(u)}u_1$.

Notice that the (potential) Burgers' equation is a particular case of (2.1). If we take $\alpha = 1$ and $\frac{\gamma - f'(u)}{f(u)} = 1$, i.e., $f(u) = \beta \exp(-u) + \gamma$, it leads to the nontrivial equation:

$$u_t = u_2 + u_1^2 + \beta \exp(-u) + \gamma.$$

2.3 Diffusion equation

Reference: [Oev84, p. 39];

Equation	$u_t = u^2 u_2$
Hamiltonian	None
Cosymplectic	None
Symplectic	None
Recursion	$u D_x + u^2 u_2 D_x^{-1} \frac{1}{u^2}$
Root	$u^2 u_2$
Scaling	$\alpha x u_1 + \beta u, \alpha, \beta \in \mathcal{C}.$

2.4 Nonlinear diffusion equation

Reference: [Olv93, Ex. 5.10];

Equation	$u_t = D_x(\frac{u_1}{u^2})$
Hamiltonian	None
Cosymplectic	None
Symplectic	None
Recursion	$D_x^2 \frac{1}{u} D_x^{-1} = \frac{1}{u} D_x - \frac{2u_1}{u^2} - u_t D_x^{-1}$
Root	u_t
Scaling	$\alpha x u_1 + \beta u, \alpha, \beta \in \mathcal{C}.$

2.5 Korteweg–de Vries equation

Reference: [Olv93, p. 312], [Oev84, pp. 18, 67, 78, 84, 97], [Dor93, pp. 85, 151, 158, 162], [Oev90, pp. 27, 60];

Equation	$u_t = u_3 + u u_1$	[KdV95]
Hamiltonian	$\frac{u^2}{2}$	
Cosymplectic	$\bar{D}_x^3 + \frac{1}{3}(u D_x + D_x u)$	
Symplectic	D_x^{-1}	
Recursion	$D_x^2 + \frac{2}{3}u + \frac{1}{3}u_1 D_x^{-1}$	[Olv77]
Root	u_1	
Scaling	$x u_1 + 2u$	

2.6 Potential Korteweg–de Vries equation

Reference: [Dor93, p. 125];

Equation	$u_t = u_3 + 3u_1^2$
Hamiltonian	$\frac{1}{2}u u_4 + 2u u_1 u_2$
Cosymplectic	D_x^{-1}
Symplectic	$D_x^3 + 2(u_1 D_x + D_x u_1)$
Recursion	$D_x^2 + 4u_1 - 2D_x^{-1}u_2$
Root	1
Scaling	$x u_1 + u$

2.7 Modified Korteweg–de Vries equation

Reference: [Olv93, Ex. 5.11], [Oev84, p. 97], [Oev90, pp. 29, 60];

Equation	$u_t = u_3 + u^2 u_1$
Hamiltonian	$\frac{u^2}{2}$
Cosymplectic	$D_x^3 + \frac{2}{3} D_x u D_x^{-1} u D_x$
Symplectic	D_x^{-1}
Recursion	$D_x^2 + \frac{2}{3} u^2 + \frac{2}{3} u_1 D_x^{-1} u$ [Olv77]
Root	u_1
Scaling	$x u_1 + u$

2.8 Potential modified Korteweg–de Vries equation

Reference: [Olv93, Ex.5.11];

Equation	$u_t = u_3 + \frac{1}{3} u_1^3$
Hamiltonian	$\frac{1}{2} u u_4 + \frac{1}{4} u u_1^2 u_2$
Cosymplectic	D_x^{-1}
Symplectic	$D_x^3 + \frac{2}{3} D_x u_1 D_x^{-1} u_1 D_x$
Recursion	$D_x^2 + \frac{2}{3} u_1^2 - \frac{2}{3} u_1 D_x^{-1} u_2$
Root	u_1
Scaling	$\alpha x u_1 + \beta u, \alpha, \beta \in \mathbb{C}$.

In the paper [Fok80], the author found the 3rd-order equations, not involving 2nd-order derivatives, i.e., of the form $u_t = u_3 + f(u, u_1)$, which possess a 5th-order symmetry and obtained the following equations:

$$u_t = u_3 + \alpha u_1^3 + \beta u_1^2 + \gamma u_1, \quad (2.3)$$

where α, β, γ are constant, with the recursion operator $D_x^2 + 2\alpha u_1^2 + \frac{4}{3}\beta u_1 - \frac{2}{3}(3\alpha u_1 + \beta)D_x^{-1}u_2 + \gamma$.

$$u_t = u_3 + \alpha u_1^3 + f(u)u_1, \quad (2.4)$$

where $f(u)$ satisfies $f''' + 8\alpha f' = 0$, with the recursion operator $D_x^2 + 2\alpha u_1^2 + \frac{2}{3}f(u) - \frac{1}{3}u_1 D_x^{-1}(6\alpha u_2 - f')$.

pkdV ($\alpha = 0$) and pmKdV ($\beta = 0$) are particular cases of (2.3). pmKdV ($f(u) = 0$), KdV ($\alpha = 0$ and $f(u) = u$) and mKdV ($\alpha = 0$ and $f(u) = u^2$) are particular cases of (2.4) including Calogero–Degasperis–Fokas equation [CD81]:

$$u_t = u_3 - \frac{1}{8} u_1^3 + (a \exp(u) + b \exp(-u) + c) u_1.$$

2.9 Cylindrical Korteweg–de Vries equation

Reference: [OF84, ZC86, Cho87a];

Equation	$u_t = u_3 + u u_1 - \frac{u}{2t}$
Hamiltonian	None
Cosymplectic	$D_x^3 + \frac{1}{3}(u D_x + D_x u) + \frac{1}{6t}(x D_x + D_x x)$
Symplectic	$t D_x^{-1}$
Recursion	$t(D_x^2 + \frac{2}{3}u + \frac{1}{3}u_1 D_x^{-1}) + \frac{1}{3}x + \frac{1}{6}D_x^{-1}$
Root	$\sqrt{t}(\frac{u_1}{3} + \frac{1}{6t})$
Scaling	$-3t\partial_t + (2u + x u_1)\partial_u$

Consider the generalized Korteweg–de Vries equation

$$u_t + u_3 + 6uu_1 + 6f(t)u - x(f'_t + 12f^2) = 0,$$

where f is an arbitrary function of t . It possesses recursion operator:

$$\mathfrak{R} = \frac{1}{g(t)}(D_x^2 + 4(u - xf(t)) + 2(u_1 - f(t))D_x^{-1})$$

with the root $\frac{1}{\sqrt{g}}(u_1 - f)$, where $g(t) = \exp(-\int 12f dt)$, cf. [Cho87a] (the author also studied generalized mKdV in the same way [Cho87b]).

If we take $f(t) = \frac{1}{12t}$ and then do transformation $\tilde{u} = \frac{1}{6}u$ and $\tilde{t} = -t$, we get the cylindrical Korteweg–de Vries equation.

2.10 Ibragimov–Shabat equation

Reference: [IŠ80, Cal87];

Equation	$u_t = u_3 + 3u^2u_2 + 9uu_1^2 + 3u^4u_1$
Hamiltonian	None
Cosymplectic	None
Symplectic	None
Root	u_1
Scaling	$xu_1 + \frac{1}{2}u$
Master Symmetries	$xu_t + \frac{3}{2}u_2 + 5u_1u^2 + \frac{1}{2}u^5$

No recursion operator seems to be known for this equation.

We should mention that this equation possesses infinitely many symmetries [IŠ80], but only one local conserved density u^2 [Kap82]. The transformation $u = \sqrt{\frac{w_1}{2w}}$ [Cal87] transforms it into $w_t = w_3 - \frac{3}{4}\frac{w_2^2}{w_1}$ and the master symmetry becomes $xw_t + \frac{1}{2}w_2$, which is the master symmetry for this new equation. Notice that the new equation has a recursion operator $\mathfrak{R} = D_x^2 - \frac{w_2}{w_1}D_x + \frac{w_3}{2w_1} - \frac{w_2^2}{4w_1^2} - D_x^{-1}(\frac{w_4}{2w_1} - \frac{w_2w_3}{w_1^2} + \frac{w_3^2}{2w_1^3}) = \left(D_x - D_x^{-1}\frac{w_2}{2w_1}D_x\right)^2$ with the root w_1 .

2.11 Harry Dym equation

Reference: [Olv93, Ex.5.15], [Oev84, p. 107];

Equation	$u_t = u^3u_3$
Hamiltonian	$-\frac{1}{u}$
Cosymplectic	$u^3D_x^3u^3$
Symplectic	$\frac{1}{u^2}D_x^{-1}\frac{1}{u^2}$
Recursion	$u^3D_x^3uD_x^{-1}\frac{1}{u^2}$ $= u^2D_x^2 - uu_1D_x + uu_2 + u^3u_3D_x^{-1}\frac{1}{u^2}$ [LLS ⁺ 83]
Root	u^3u_3
Scaling	$\alpha xu_1 + \beta u, \alpha, \beta \in \mathbb{C}$.

Sometimes the equation is written as $u_t = D_x^3(\frac{1}{\sqrt{u}})$, cf. [Dor93, p. 85].

2.12 Schwarzian KdV equation

Reference: [Dor93, p. 121];

Equation	$u_t = u_3 - \frac{3}{2} \frac{u_2^2}{u_1}$
Hamiltonian	$\frac{u_2^2}{2u_1^2}$
Cosymplectic	$2(\frac{1}{u_1^2} D_x + D_x \frac{1}{u_1^2})^{-1}$
Symplectic	$\frac{1}{2}(\frac{1}{u_1^2} D_x^3 + D_x^3 \frac{1}{u_1^2}) + (\frac{u_3}{u_1^3} - \frac{3u_2^2}{u_1^4}) D_x + D_x (\frac{u_3}{u_1^3} - \frac{3u_2^2}{u_1^4})$
Recursion	$\begin{cases} D_x^2 - \frac{2u_2}{u_1} D_x + (\frac{u_3}{u_1^3} - \frac{u_2^2}{u_1^4}) - u_1 D_x^{-1} \xi, \\ \xi = \frac{3u_2^3}{u_1^4} - \frac{4u_2 u_3}{u_1^3} + \frac{u_4}{u_1^2} \end{cases}$
Root	u_1
Scaling	$\alpha x u_1 + \beta u, \alpha, \beta \in \mathbb{C}.$

2.13 Cavalcante–Tenenblat equation

Reference: [CT88];

Equation	$u_t = D_x^2(u_1^{-\frac{1}{2}}) + u_1^{\frac{3}{2}}$
Hamiltonian	$-2\sqrt{u_1}$
Cosymplectic	$D_x - u_1 D_x^{-1} u_1$
Symplectic	$u_1^{-\frac{1}{2}} D_x u_1^{-\frac{1}{2}} - \frac{1}{4} u_1^{-\frac{3}{2}} u_2 D_x^{-1} u_1^{-\frac{3}{2}} u_2$
Recursion	$\frac{1}{u_1} D_x^2 - \frac{3u_2}{2u_1^2} D_x - \frac{u_3}{2u_1^3} + \frac{3u_2^2}{4u_1^3} - u_1 + \frac{u_t}{2} D_x^{-1} u_1^{-\frac{3}{2}} u_2$
Root	u_t
Scaling	$x u_1$

2.14 Liouville equation

Reference: [Dor93, pp. 134, 164];

Equation	$u_{xt} = \exp(u)$	
Hamiltonian	$\exp(u)$	★
Cosymplectic	D_x^{-1}	
Symplectic	$D_x^3 - D_x u_1 D_x^{-1} u_1 D_x$	
Recursion	$D_x^2 - u_1^2 + u_1 D_x^{-1} u_2$	
Root	u_1	
Scaling	$\alpha x u_1 + \beta u, \alpha, \beta \in \mathbb{C}.$	

★ Actually, the equation is treated as an evolution equation $u_t = D_x^{-1} \exp(u)$.

The **Sinh–Gordon equation** $u_{xt} = \sinh u$ has exactly the same geometric structure, cf. [AC91].

We refer to [ZS01] for recent developments in integrable hyperbolic equations of Liouville type.

2.15 Sine–Gordon equation

Reference: [Olv93, Ex.5.12], [Dor93, pp. 133, 163];

Equation	$u_{xt} = \sin u$	Oevel[16]
Hamiltonian	$-\cos u$	\star
Cosymplectic	D_x^{-1}	
Symplectic	$D_x^3 + D_x u_1 D_x^{-1} u_1 D_x$	
Recursion	$D_x^2 + u_1^2 - u_1 D_x^{-1} u_2$	[Olv77]
Root	u_1	
Scaling	$\alpha x u_1 + \beta u, \alpha, \beta \in \mathbb{C}$	

\star As we mentioned for the Liouville equation, this equation is also treated as an evolution equation $u_t = D_x^{-1} \sin u$.

2.16 Klein–Gordon equations

Reference: [AC91, p. 366], [Kon87, p. 41], [FG80];

Equation	$u_{xt} = \alpha \exp(-2u) + \beta \exp(u)$
Hamiltonian	$-\frac{\alpha}{2} \exp(-2u) + \beta \exp(u)$
Cosymplectic	D_x^{-1}
Symplectic	\mathfrak{J}
Recursion	\mathfrak{R}
Root	$u_1, u_5 + 5u_2u_3 - 5u_1^2u_3 - 5u_1u_2^2 + u_1^5$
Scaling	xu_1

$$\begin{aligned}
\mathfrak{J} &= D_x^7 + 3(u_2 D_x^5 + D_x^5 u_2) - 3(u_1^2 D_x^5 + D_x^5 u_1^2) - 8(u_4 D_x^3 + D_x^3 u_4) \\
&\quad + 10(u_1 u_3 D_x^3 + D_x^3 u_1 u_3) + \frac{29}{2}(u_2^2 D_x^3 + D_x^3 u_2^2) - 3(u_1^2 u_2 D_x^3 + D_x^3 u_1^2 u_2) \\
&\quad + \frac{9}{2}(u_1^4 D_x^3 + D_x^3 u_1^4) + 5(u_6 D_x + D_x u_6) - 6(u_1 u_5 D_x + D_x u_1 u_5) \\
&\quad - 25(u_2 u_4 D_x + D_x u_2 u_4) + 3(u_1^2 u_4 D_x + D_x u_1^2 u_4) - 21(u_3^2 D_x + D_x u_3^2) \\
&\quad + 8(u_1 u_2 u_3 D_x + D_x u_1 u_2 u_3) - 8(u_1^3 u_3 D_x + D_x u_1^3 u_3) \\
&\quad + 6(u_2^3 D_x + D_x u_2^3) - 44(u_1^2 u_2^2 D_x + D_x u_1^2 u_2^2) - 2(u_1^6 D_x + D_x u_1^6) \\
&\quad + 2u_2 D_x^{-1}(u_6 + 5u_2 u_4 + 5u_3^2 - 5u_1^2 u_4 - 20u_1 u_2 u_3 - 5u_2^3 + 5u_1^4 u_2) \\
&\quad + 2(u_6 + 5u_2 u_4 + 5u_3^2 - 5u_1^2 u_4 - 20u_1 u_2 u_3 - 5u_2^3 + 5u_1^4 u_2) D_x^{-1} u_2 \\
\mathfrak{R} &= D_x^6 + 6(u_2 - u_1^2) D_x^4 + 9(u_3 - 2u_1 u_2) D_x^3 \\
&\quad + (5u_4 - 22u_1 u_3 - 13u_2^2 - 6u_1^2 u_2 + 9u_1^4) D_x^2 \\
&\quad + (u_5 - 8u_1 u_4 - 15u_2 u_3 - 3u_1^2 u_3 - 6u_1 u_2^2 + 18u_1^3 u_2) D_x \\
&\quad - 4u_1 u_5 + 20u_1^3 u_3 - 20u_1 u_2 u_3 + 20u_1^2 u_2^2 - 4u_1^6 \\
&\quad + 2u_1 D_x^{-1}(u_6 + 5u_2 u_4 + 5u_3^2 - 5u_1^2 u_4 - 20u_1 u_2 u_3 - 5u_2^3 + 5u_1^4 u_2) \\
&\quad + 2(u_5 + 5u_2 u_3 - 5u_1^2 u_3 - 5u_1 u_2^2 + u_1^5) D_x^{-1} u_2
\end{aligned}$$

It shares its recursion operator [Bil93] with the Potential Kupershmidt equation, i.e., $u_t = u_5 + 5u_2 u_3 - 5u_1^2 u_3 - 5u_1 u_2^2 + u_1^5$ (equation (4.2.7) in [MSS91]).

Klein–Gordon equations $u_{xt} = f(u)$ possess a nontrivial symmetry if and only if $f(u) = \alpha \exp(-\lambda u) + \beta \exp(\lambda u)$ or $f(u) = \alpha \exp(-2\lambda u) + \beta \exp(\lambda u)$, cf. [ZS79].

2.17 Kupershmidt equation

Reference: [MSS91, Eq. (4.2.6)], [FG80, Bil93];

Equation	$u_t = u_5 + 5u_1u_3 + 5u_2^2 - 5u^2u_3 - 20uu_1u_2 - 5u_1^3 + 5u^4u_1$
Hamiltonian	$\frac{u_2^2}{2} - \frac{5u_1^3}{6} + \frac{5u^2u_1^2}{2} + \frac{u^6}{6}$
Cosymplectic	\bar{D}_x
Symplectic	\mathfrak{I}
Recursion	\mathfrak{R}
Root	u_1, u_t
Scaling	$xu_1 + u$
\mathfrak{I}	$D_x^5 + 3(u_1D_x^3 + D_x^3u_1) - 3(u^2D_x^3 + D_x^3u^2) - 3(u_1u^2D_x + D_xu_1u^2)$ $+ \frac{5}{2}(u_1^2D_x + D_xu_1^2) - 2(u_3D_x + D_xu_3) + \frac{9}{2}(u^4D_x + D_xu^4)$ $- 2(uu_2D_x + D_xuu_2) - 2(u_4 - 5u^2u_2 - 5uu_1^2 + 5u_1u_2 + u^5)D_x^{-1}u$ $- 2uD_x^{-1}(u_4 - 5u^2u_2 - 5uu_1^2 + 5u_1u_2 + u^5)$
\mathfrak{R}	$D_x^6 + 6u_1D_x^4 - 6u^2D_x^4 - 30uu_1D_x^3 + 15u_2D_x^3 + 9u^4D_x^2 - 6u^2u_1D_x^2$ $- 40uu_2D_x^2 - 31u_1^2D_x^2 + 14u_3D_x^2 - 9u^2u_2D_x + 54u^3u_1D_x - 18uu_1^2D_x$ $- 30uu_3D_x - 63u_1u_2D_x + 6u_4D_x - 4u^6 + 38u^3u_2 + 74u^2u_1^2$ $- 3u^2u_3 - 12uu_4 - 38uu_1u_2 + u_5 - 6u_1^3 - 23u_1u_3 - 15u_2^2$ $- 2u_tD_x^{-1}u - 2u_1D_x^{-1}(u_4 - 5u^2u_2 - 5uu_1^2 + 5u_1u_2 + u^5)$

2.18 Sawada–Kotera equation

Reference: [SK74, CDG76, FO82, FOW87, Bil93], [Oev84, p. 105], [Oev90, p. 30], [MSS91, Eq. (4.2.2)];

Equation	$u_t = u_5 + 5uu_3 + 5u_1u_2 + 5u^2u_1$
Hamiltonian	$\frac{u^3}{6} - \frac{u_1^2}{2}$
Cosymplectic	$D_x (D_x + 2(D_x^{-1}u + uD_x^{-1})) D_x$
Symplectic	$(D_x + D_x^{-1}u) D_x (D_x + uD_x^{-1})$
Recursion	\mathfrak{R}
Root	u_1, u_t
Scaling	$xu_1 + 2u$
\mathfrak{R}	$D_x^6 + 6uD_x^4 + 9u_1D_x^3 + 9u^2D_x^2 + 11u_2D_x^2 + 10u_3D_x + 21uu_1D_x$ $+ 4u^3 + 16uu_2 + 6u_1^2 + 5u_4 + u_1D_x^{-1}(2u_2 + u^2) + u_tD_x^{-1}$

2.19 Potential Sawada–Kotera equation

Reference: [MSS91, Eq. (4.2.4)], [Bil93];

Equation	$u_t = u_5 + 5u_1u_3 + \frac{5}{3}u_1^3$
Hamiltonian	$\frac{u_2^2}{2} - \frac{u_1^3}{6}$
Cosymplectic	$D_x + 2(u_1D_x^{-1} + D_x^{-1}u_1)$
Symplectic	$(D_x + u_1D_x^{-1}) D_x^3 (D_x + D_x^{-1}u_1)$
Recursion	\mathfrak{R}
Root	$u_1, 1$
Scaling	$xu_1 + u$
\mathfrak{R}	$D_x^6 + 6u_1D_x^4 + 3u_2D_x^3 + 8u_3D_x^2 + 9u_1^2D_x^2 + 2u_4D_x + 3u_2u_1D_x$

$$+3u_5 + 13u_3u_1 + 3u_2^2 + 4u_1^3 - 2u_1D_x^{-1}(u_4 + u_2u_1) \\ -2D_x^{-1}(u_6 + 3u_4u_1 + 6u_3u_2 + 2u_2u_1^2)$$

2.20 Kaup–Kupershmidt equation

Reference: [Kau80, FO82, FOW87, Bil93], [MSS91, Eq. (4.2.3)];

Equation	$u_t = u_5 + 5uu_3 + \frac{25}{2}u_1u_2 + 5u^2u_1$
Hamiltonian	$\frac{2u^3}{3} - \frac{u_1^2}{2}$
Cosymplectic	$D_x \left(D_x + \frac{1}{2}(uD_x^{-1} + D_x^{-1}u) \right) D_x$
Symplectic	$D_x^3 + \frac{3}{2}(uD_x + D_xu) + D_x^2uD_x^{-1} + D_x^{-1}uD_x^2 \\ + 2(u^2D_x^{-1} + D_x^{-1}u^2)$
Recursion	\mathfrak{R}
Root	u_1, u_t
Scaling	$xu_1 + 2u$

$$\mathfrak{R} = D_x^6 + 6uD_x^4 + 18u_1D_x^3 + 9u^2D_x^2 + \frac{49}{2}u_2D_x^2 + 30uu_1D_x + \frac{35}{2}u_3D_x \\ + 4u^3 + \frac{41}{2}uu_2 + \frac{69}{4}u_1^2 + \frac{13}{2}u_4 + \frac{1}{2}u_1D_x^{-1}(u_2 + 2u^2) + u_tD_x^{-1}$$

2.21 Potential Kaup–Kupershmidt equation

Reference: [MSS91, Eq. (4.2.5)], [Bil93];

Equation	$u_t = u_5 + 10u_1u_3 + \frac{15}{2}u_2^2 + \frac{20}{3}u_1^3$
Hamiltonian	$\frac{u_2^2}{2} - \frac{4u_1^3}{3}$
Cosymplectic	$D_x + u_1D_x^{-1} + D_x^{-1}u_1$
Symplectic	$D_x^5 + 5(u_1D_x^3 + D_x^3u_1) - 3(u_3D_x + D_xu_3) \\ + 8(u_1^2D_x + D_xu_1^2)$
Recursion	\mathfrak{R}
Root	$u_1, 1$
Scaling	$xu_1 + u$

$$\mathfrak{R} = D_x^6 + 12u_1D_x^4 + 24u_2D_x^3 + 25u_3D_x^2 + 36u_1^2D_x^2 + 10u_4D_x + 48u_1u_2D_x \\ + 3u_5 + 21u_2^2 + 34u_1u_3 + 32u_1^3 - 2u_1D_x^{-1}(u_4 + 8u_1u_2) \\ - D_x^{-1}(u_6 + 12u_1u_4 + 24u_2u_3 + 32u_1^2u_2)$$

2.22 Diffusion system

Reference: [Oev84, p. 41];

Equation	$\begin{cases} u_t = u_2 + v^2 \\ v_t = v_2 \end{cases}$
Hamiltonian	None
Cosymplectic	None
Symplectic	None
Recursion	$\begin{pmatrix} D_x & vD_x^{-1} \\ 0 & D_x \end{pmatrix}$
Root	$\begin{pmatrix} v \\ 0 \end{pmatrix}, \begin{pmatrix} u_1 \\ v_1 \end{pmatrix}$

$$\text{Scaling} \quad \begin{pmatrix} xu_1 + 2u \\ xv_1 + 2v \end{pmatrix} + \alpha \begin{pmatrix} 2u \\ v \end{pmatrix}, \alpha \in \mathbb{C}$$

2.23 Dispersiveless Long Wave system

Reference: [AC91, Gök98];

$$\begin{array}{ll} \text{Equation} & \begin{cases} u_t = u_1 v + uv_1 \\ v_t = u_1 + vv_1 \end{cases} \\ \text{Hamiltonian} & \frac{u^2 + uv^2}{2} \\ \text{Cosymplectic} & \begin{pmatrix} 0 & D_x \\ D_x & 0 \end{pmatrix} \\ \text{Symplectic} & \begin{pmatrix} 2D_x^{-1} & vD_x^{-1} \\ D_x^{-1}v & uD_x^{-1} + D_x^{-1}u \end{pmatrix} \\ \text{Recursion} & \begin{pmatrix} v & 2u + u_1 D_x^{-1} \\ 2 & v + v_1 D_x^{-1} \end{pmatrix} \\ \text{Root} & \begin{pmatrix} u_1 \\ v_1 \end{pmatrix} \\ \text{Scaling} & \begin{pmatrix} xu_1 \\ xv_1 \end{pmatrix} + \alpha \begin{pmatrix} 2u \\ v \end{pmatrix}, \alpha \in \mathbb{C} \end{array}$$

2.24 Sine-Gordon equation in the laboratory coordinates

Reference: [CLL87];

$$\begin{array}{ll} \text{Equation} & \begin{cases} u_t = v \\ v_t = u_2 - \sin(u) \end{cases} \\ \text{Hamiltonian} & \frac{1}{2}(u_1^2 + v^2) - \cos(u) \\ \text{Cosymplectic} & \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \\ \text{Symplectic} & \begin{pmatrix} -\mathfrak{R}_{21} & -\mathfrak{R}_{22} \\ \mathfrak{R}_{11} & \mathfrak{R}_{12} \end{pmatrix} \\ \text{Recursion} & \mathfrak{R} = \begin{pmatrix} \mathfrak{R}_{11} & \mathfrak{R}_{12} \\ \mathfrak{R}_{21} & \mathfrak{R}_{22} \end{pmatrix} \\ \text{Root} & \begin{pmatrix} u_1 \\ v_1 \end{pmatrix} \\ \text{Scaling} & \text{None} \end{array}$$

$$\begin{aligned} \mathfrak{R}_{11} &= 4D_x^2 - 2\cos(u) + (u_1 + v)^2 - (u_1 + v)D_x^{-1}(u_2 + v_1 - \sin(u)), \\ \mathfrak{R}_{12} &= 4D_x + (u_1 + v)D_x^{-1}(u_1 + v), \\ \mathfrak{R}_{21} &= 4D_x^3 + (u_1 + v)^2 D_x - 4\cos(u)D_x + 2u_1 \sin(u) + (u_2 + v_1)(u_1 + v) \\ &\quad - (u_2 + v_1 - \sin(u))D_x^{-1}(u_2 + v_1 - \sin(u)), \\ \mathfrak{R}_{22} &= 4D_x^2 + (u_1 + v)^2 - 2\cos(u) + (u_2 + v_1 - \sin(u))D_x^{-1}(u_1 + v). \end{aligned}$$

2.25 AKNS equation

Reference: [Oev84, p. 100];

Equation	$\begin{cases} u_t = -u_2 + 2u^2v \\ v_t = v_2 - 2v^2u \end{cases}$
Hamiltonian	$\frac{1}{2}(uv_1 - vu_1)$
Cosymplectic	$\begin{pmatrix} 2uD_x^{-1}u & D_x - 2uD_x^{-1}v \\ D_x - 2vD_x^{-1}u & 2vD_x^{-1}v \end{pmatrix}$
Symplectic	$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$
Recursion	$\begin{pmatrix} -D_x + 2uD_x^{-1}v & 2uD_x^{-1}u \\ -2vD_x^{-1}v & D_x - 2vD_x^{-1}u \end{pmatrix}$
Root	$\begin{pmatrix} -u \\ v \end{pmatrix}$
Scaling	$\begin{pmatrix} xu_1 + u \\ xv_1 + v \end{pmatrix}$

2.26 Nonlinear Schrödinger equation

Reference: [Oev84, p. 102], [Dor93, p. 135], [Oev90, pp. 31, 61];

Equation	$\begin{cases} u_t = v_2 \mp v(u^2 + v^2) \\ v_t = -u_2 \pm u(u^2 + v^2) \end{cases}$
Hamiltonian	$\frac{1}{2}(uv_1 - vu_1)$
Cosymplectic	$\begin{pmatrix} D_x \mp 2vD_x^{-1}v & \pm 2vD_x^{-1}u \\ \pm 2uD_x^{-1}v & D_x \mp 2uD_x^{-1}u \end{pmatrix}$
Symplectic	$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$
Recursion	$\begin{pmatrix} \mp 2vD_x^{-1}u & D_x \mp 2vD_x^{-1}v \\ -D_x \pm 2uD_x^{-1}u & \pm 2uD_x^{-1}v \end{pmatrix}$
Root	$\begin{pmatrix} -v \\ u \end{pmatrix}$
Scaling	$\begin{pmatrix} xu_1 + u \\ xv_1 + v \end{pmatrix}$

The system can be written as $iq_t = q_2 \mp q^2 q^*$, where $i^2 = -1$, cf. [AC91].

2.27 Derivative Schrödinger system

Reference: [Oev84, p. 103];

Equation	$\begin{cases} u_t = -v_2 - (u^2 + v^2)u_1 \\ v_t = u_2 - (u^2 + v^2)v_1 \end{cases}$
Hamiltonian	$\frac{1}{2}(uv_1 - vu_1)$
Cosymplectic	$\begin{pmatrix} -D_x & \frac{u^2+v^2}{2} \\ -\frac{u^2+v^2}{2} & -D_x \end{pmatrix} - \begin{pmatrix} v \\ -u \end{pmatrix} D_x^{-1} \begin{pmatrix} u_1 \\ v_1 \end{pmatrix} - \begin{pmatrix} u_1 \\ v_1 \end{pmatrix} D_x^{-1} \begin{pmatrix} v \\ -u \end{pmatrix}$
Symplectic	$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$

$$\begin{array}{ll}
\text{Recursion} & \begin{pmatrix} -\frac{u^2+v^2}{2} & -D_x \\ D_x & -\frac{u^2+v^2}{2} \end{pmatrix} + \begin{pmatrix} v \\ -u \end{pmatrix} D_x^{-1} \begin{pmatrix} v_1 \\ -u_1 \end{pmatrix}^\dagger \\
& - \begin{pmatrix} u_1 \\ v_1 \end{pmatrix} D_x^{-1} \begin{pmatrix} u \\ v \end{pmatrix}^\dagger \\
\text{Root} & \begin{pmatrix} v \\ -u \end{pmatrix} \\
\text{Scaling} & \begin{pmatrix} xu_1 + \frac{u}{2} \\ xv_1 + \frac{v}{2} \end{pmatrix}
\end{array}$$

2.28 Modified derivative Schrödinger system

Reference: [WHV95];

$$\begin{array}{ll}
\text{Equation} & \begin{cases} u_t = D_x(u^3 + uv^2 + \beta u - v_1) \\ v_t = D_x(vu^2 + v^3 + u_1) \end{cases} \\
\text{Hamiltonian} & \frac{1}{2}(u^2 + v^2) \\
\text{Cosymplectic} & \begin{pmatrix} \beta D_x + 2uD_x u & -D_x^2 + 2vD_x u \\ D_x^2 + 2uD_x v & 2vD_x v \end{pmatrix} \\
& -2 \begin{pmatrix} u_1 \\ v_1 \end{pmatrix} D_x^{-1} \begin{pmatrix} u_1 \\ v_1 \end{pmatrix} \\
\text{Symplectic} & \begin{pmatrix} D_x^{-1} & 0 \\ 0 & D_x^{-1} \end{pmatrix} \\
\text{Recursion} & \begin{pmatrix} \beta + 2u^2 & -D_x + 2uv \\ D_x + 2uv & 2v^2 \end{pmatrix} \\
& +2 \begin{pmatrix} u_1 \\ v_1 \end{pmatrix} D_x^{-1} \begin{pmatrix} u \\ v \end{pmatrix}^\dagger \\
\text{Root} & \begin{pmatrix} u_1 \\ v_1 \end{pmatrix} \\
\text{Scaling} & \begin{pmatrix} xu_1 + \frac{u}{2} \\ xv_1 + \frac{v}{2} \\ x\beta_x + \beta \end{pmatrix}.
\end{array}$$

2.29 Boussinesq system

Reference: [Olv93, p. 459];

$$\begin{array}{ll}
\text{Equation} & \begin{cases} u_t = v_1 \\ v_t = \frac{1}{3}u_3 + \frac{8}{3}uu_1 \end{cases} \\
\text{Hamiltonian} & \frac{1}{2}v \\
\text{Cosymplectic} & \begin{pmatrix} D_x^3 + 2uD_x + u_1 & 3vD_x + 2v_1 \\ 3vD_x + v_1 & \mathfrak{H}_{22} \end{pmatrix} \\
\text{Symplectic} & \begin{pmatrix} 0 & D_x^{-1} \\ D_x^{-1} & 0 \end{pmatrix} \\
\text{Recursion} & \begin{pmatrix} 3v + 2v_1 D_x^{-1} & D_x^2 + 2u + u_1 D_x^{-1} \\ \mathfrak{R}_{21} & 3v + v_1 D_x^{-1} \end{pmatrix} \\
\text{Root} & \begin{pmatrix} u_1 \\ v_1 \end{pmatrix}, \begin{pmatrix} u_t \\ v_t \end{pmatrix}
\end{array}$$

$$\text{Scaling} \quad \begin{pmatrix} xu_1 + 2u \\ xv_1 + 3v \end{pmatrix}$$

$$\begin{aligned} \mathfrak{H}_{22} &= \frac{1}{3}D_x^5 + \frac{5}{3}(uD_x^3 + D_x^3u) - (u_2D_x + D_xu_2) + \frac{16}{3}uD_xu \\ \mathfrak{R}_{21} &= \frac{1}{3}D_x^4 + \frac{10}{3}uD_x^2 + 5u_1D_x + 3u_2 + \frac{16}{3}u^2 + 2v_tD_x^{-1} \end{aligned}$$

2.30 Modified Boussinesq system

Reference: [FG81];

$$\begin{aligned} \text{Equation} & \quad \begin{cases} u_t = 3v_2 + 6uv_1 + 6u_1v \\ v_t = -u_2 - 6vv_1 + 2uu_1 \end{cases} \\ \text{Hamiltonian} & \quad \frac{1}{2}(uv_1 - u_1v - 2v^3 + 2vu^2) \\ \text{Cosymplectic} & \quad \begin{pmatrix} 3D_x & 0 \\ 0 & D_x \end{pmatrix} \\ \text{Symplectic} & \quad \begin{pmatrix} \frac{1}{3}D_x^{-1}\mathfrak{R}_{11} & \frac{1}{3}D_x^{-1}\mathfrak{R}_{12} \\ D_x^{-1}\mathfrak{R}_{21} & D_x^{-1}\mathfrak{R}_{22} \end{pmatrix} \\ \text{Recursion} & \quad \begin{pmatrix} \mathfrak{R}_{11} & \mathfrak{R}_{12} \\ \mathfrak{R}_{21} & \mathfrak{R}_{22} \end{pmatrix} \\ \text{Root} & \quad \begin{pmatrix} u_1 \\ v_1 \end{pmatrix}, \begin{pmatrix} u_t \\ v_t \end{pmatrix} \\ \text{Scaling} & \quad \begin{pmatrix} xu_1 + u \\ xv_1 + v \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \mathfrak{R}_{11} &= 6vD_x^2 + 9v_1D_x + 3v_2 - 12uv_1 - 24u^2v - 2u_tD_x^{-1}u - 6u_1D_x^{-1}(2uv + v_1), \\ \mathfrak{R}_{12} &= 3D_x^3 + 6uD_x^2 + 9u_1D_x - 3u^2D_x - 9v^2D_x + 3u_2 - 6u^3 - 36vv_1 \\ &\quad - 18uv^2 - 6u_tD_x^{-1}v + 6u_1D_x^{-1}(u_1 - u^2 + 3v^2), \\ \mathfrak{R}_{21} &= -D_x^3 + 2uD_x^2 + u^2D_x + 3u_1D_x + 3v^2D_x + u_2 - 6uv^2 - 2u^3 + 4uu_1 \\ &\quad - 2v_tD_x^{-1}u - 6v_1D_x^{-1}(v_1 + 2uv), \\ \mathfrak{R}_{22} &= -6vD_x^2 - 9v_1D_x - 12u^2v + 12u_1v - 3v_2 + 36v^3 \\ &\quad - 6v_tD_x^{-1}v + 6v_1D_x^{-1}(u_1 - u^2 + 3v^2). \end{aligned}$$

2.31 Landau–Lifshitz system

Reference: [vBK91];

$$\begin{aligned} \text{Equation} & \quad \begin{cases} u_t = -\sin(u)v_2 - 2\cos(u)u_1v_1 + (J_1 - J_2)\sin(u)\cos(v)\sin(v) \\ v_t = \frac{u_2}{\sin(u)} - \cos(u)v_1^2 + \cos(u)(J_1\cos^2(v) + J_2\sin^2(v) - J_3) \end{cases} \\ \text{Hamiltonian} & \quad \frac{1}{2}(\sin^2(u)(J_1\cos^2(v) + J_2\sin^2(v) - J_3) + J_3 - u_1^2 - \sin^2(u)v_1^2) \\ \text{Cosymplectic} & \quad \frac{1}{\sin(u)} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \\ \text{Symplectic} & \quad \sin(u) \begin{pmatrix} \mathfrak{R}_{21} & \mathfrak{R}_{22} \\ -\mathfrak{R}_{11} & -\mathfrak{R}_{12} \end{pmatrix} \end{aligned}$$

$$\begin{array}{ll}
\text{Recursion} & \begin{pmatrix} \mathfrak{R}_{11} & \mathfrak{R}_{12} \\ \mathfrak{R}_{21} & \mathfrak{R}_{22} \end{pmatrix} \\
\text{Root} & \begin{pmatrix} u_1 \\ v_1 \end{pmatrix}, \begin{pmatrix} u_t \\ v_t \end{pmatrix} \\
\text{Scaling} & \begin{pmatrix} xu_1 \\ xv_1 \\ xJ_x + 2J \end{pmatrix}, \text{ where } J = (J_1, J_2, J_3).
\end{array}$$

where

$$\begin{aligned}
\mathfrak{R}_{11} &= -D_x^2 - 2\sin^2(u)v_1^2 - u_1^2 + v_1^2 - (J_1 - J_2)\sin^2(u)\sin^2(v) \\
&\quad + (J_1 - J_3)\sin^2(u) + J_3 - J_2 + u_tD_x^{-1} \cdot (\sin(u)v_1) - u_1D_x^{-1} \cdot S_1 \\
\mathfrak{R}_{12} &= 2\cos(u)\sin(u)v_1D_x + \cos(u)\sin(u)v_2 - 3\sin^2(u)u_1v_1 + 2u_1v_1 \\
&\quad + u_tD_x^{-1} \cdot (-\sin(u)u_1) - u_1D_x^{-1} \cdot S_2 \\
\mathfrak{R}_{21} &= -2\cos(u)v_1D_x - \cos(u)v_2 + u_1v_1 \\
&\quad + v_tD_x^{-1} \cdot (\sin(u)v_1) - v_1D_x^{-1} \cdot S_1 \\
\mathfrak{R}_{22} &= -D_x^2 - 2\cos(u)u_1D_x - \cos(u)u_2 - (J_1 - J_2)\sin(u)\sin^2(v) \\
&\quad - 2\sin^2(u)v_1^2 + v_1^2 + (J_1 - J_3)\sin^2(u) + J_3 - J_2 \\
&\quad + v_tD_x^{-1} \cdot (-\sin(u)u_1) - v_1D_x^{-1} \cdot S_2 \\
S_1 &= (J_1 - J_2)\cos(u)\sin(u)\sin^2(v) - (J_1 - J_3)\cos(u)\sin(u) \\
&\quad + \cos(u)\sin(u)v_1^2 - u_2, \\
S_2 &= (J_1 - J_2)\cos(v)\sin^2(u)\sin(v) - 2\cos(u)\sin(u)u_1v_1 - \sin^2(u)v_2.
\end{aligned}$$

2.32 Wadati–Konno–Ichikawa system

Reference: [WKI79, BPT83], [Kon87, p. 88];

$$\begin{array}{ll}
\text{Equation} & \begin{cases} u_t = D_x^2\left(\frac{u}{\sqrt{1+uv}}\right) \\ v_t = -D_x^2\left(\frac{v}{\sqrt{1+uv}}\right) \end{cases} \\
\text{Hamiltonian} & 2\sqrt{1+uv} \\
\text{Cosymplectic} & \begin{pmatrix} 0 & D_x^2 \\ -D_x^2 & 0 \end{pmatrix} \\
\text{Symplectic} & \begin{pmatrix} 0 & \frac{2}{1+uv} \\ -\frac{2}{1+uv} & 0 \end{pmatrix} - \begin{pmatrix} \frac{v}{\sqrt{1+uv}} \\ \frac{u}{\sqrt{1+uv}} \end{pmatrix}^\dagger D_x^{-1} \begin{pmatrix} \frac{v_1}{(1+uv)^{\frac{3}{2}}} \\ -\frac{u_1}{(1+uv)^{\frac{3}{2}}} \end{pmatrix}^\dagger \\
& - \begin{pmatrix} \frac{v_1}{(1+uv)^{\frac{3}{2}}} \\ -\frac{u_1}{(1+uv)^{\frac{3}{2}}} \end{pmatrix}^\dagger D_x^{-1} \begin{pmatrix} \frac{v}{\sqrt{1+uv}} \\ \frac{u}{\sqrt{1+uv}} \end{pmatrix}^\dagger \\
\text{Root} & \begin{pmatrix} u_t \\ v_t \end{pmatrix}, \begin{pmatrix} D_x^2\left(\frac{u_1}{(1+uv)^{\frac{3}{2}}}\right) \\ D_x^2\left(\frac{v_1}{(1+uv)^{\frac{3}{2}}}\right) \end{pmatrix} \\
\text{Scaling} & \begin{pmatrix} xu_1 \\ xv_1 \end{pmatrix}
\end{array}$$

2.33 Hirota–Satsuma system

Reference: [HS81, Fuc82], [Kon87, p. 207], [Oev90, pp. 32, 61], [Oev84, pp. 31, 84];

$$\begin{array}{ll}
\text{Equation} & \begin{cases} u_t = \frac{1}{2}u_3 + 3uu_1 - 6vv_1 \\ v_t = -v_3 - 3uv_1 \end{cases} \\
\text{Hamiltonian} & \frac{1}{2}u^2 - v^2 \\
\text{Cosymplectic} & \begin{pmatrix} \frac{1}{2}D_x^3 + uD_x + D_xu & vD_x + D_xv \\ vD_x + D_xv & \frac{1}{2}D_x^3 + uD_x + D_xu \end{pmatrix} \\
\text{Symplectic} & \begin{pmatrix} \frac{1}{2}D_x + uD_x^{-1} + D_x^{-1}u & -2D_x^{-1}v \\ -2vD_x^{-1} & -2D_x \end{pmatrix} \\
\text{Recursion} & \mathfrak{R} \\
\text{Root} & \begin{pmatrix} u_1 \\ v_1 \end{pmatrix}, \begin{pmatrix} u_t \\ v_t \end{pmatrix} \\
\text{Scaling} & \begin{pmatrix} xu_1 + 2u \\ xv_1 + 2v \end{pmatrix} \\
\mathfrak{R}(u, v) & = \begin{pmatrix} \frac{1}{2}D_x^3 + D_x \cdot u + uD_x & D_x \cdot v + vD_x \\ D_x \cdot v + vD_x & \frac{1}{2}D_x^3 + D_x \cdot u + uD_x \end{pmatrix} \\
& \begin{pmatrix} \frac{1}{2}D_x + D_x^{-1} \cdot u + uD_x^{-1} & -2D_x^{-1} \cdot v \\ -2vD_x^{-1} & -2D_x \end{pmatrix} \\
& \sim \begin{pmatrix} \frac{1}{2}u_3 + 3uu_1 - 6vv_1 \\ -v_3 - 3uv_1 \end{pmatrix} \otimes D_x^{-1} \begin{pmatrix} 1 & 0 \end{pmatrix} \\
& + \begin{pmatrix} u_1 \\ v_1 \end{pmatrix} \otimes D_x^{-1} \begin{pmatrix} u & -2v \end{pmatrix}
\end{array}$$

2.34 The Symmetrically-coupled Korteweg-de Vries system

Reference: [Fuc82];

$$\begin{array}{ll}
\text{Equation} & \begin{cases} u_t = u_3 + v_3 + 6uu_1 + 4uv_1 + 2u_1v \\ v_t = u_3 + v_3 + 6vv_1 + 4vu_1 + 2v_1u \end{cases} \\
\text{Hamiltonian} & \frac{1}{2}(u + v)^2 \\
\text{Cosymplectic} & \begin{pmatrix} D_x^3 + 2(uD_x + D_xu) & 0 \\ 0 & D_x^3 + 2(vD_x + D_xv) \end{pmatrix} \\
\text{Symplectic} & \begin{pmatrix} D_x^{-1} & D_x^{-1} \\ D_x^{-1} & D_x^{-1} \end{pmatrix} \\
\text{Recursion} & \begin{pmatrix} D_x^2 + 4u + 2u_1D_x^{-1} & D_x^2 + 4u + 2u_1D_x^{-1} \\ D_x^2 + 4v + 2v_1D_x^{-1} & D_x^2 + 4v + 2v_1D_x^{-1} \end{pmatrix} \\
\text{Root} & \begin{pmatrix} u_1 \\ v_1 \end{pmatrix} \\
\text{Scaling} & \begin{pmatrix} xu_1 + 2u \\ xv_1 + 2v \end{pmatrix}
\end{array}$$

2.35 The Complexly-coupled Korteweg-de Vries system

Reference: [Fuc82];

$$\begin{array}{ll}
\text{Equation} & \begin{cases} u_t = u_3 + 6uu_1 + 6vv_1 \\ v_t = v_3 + 6uv_1 + 6vu_1 \end{cases} \\
\text{Hamiltonian} & \frac{1}{2}(u^2 + v^2) \\
\text{Cosymplectic} & \begin{pmatrix} D_x^3 + 2(uD_x + D_xu) & 2D_xv + 2vD_x \\ 2D_xv + 2vD_x & D_x^3 + 2(uD_x + D_xu) \end{pmatrix}
\end{array}$$

$$\begin{array}{ll}
\text{Symplectic} & \begin{pmatrix} D_x^{-1} & 0 \\ 0 & D_x^{-1} \end{pmatrix} \\
\text{Recursion} & \begin{pmatrix} D_x^2 + 4u + 2u_1 D_x^{-1} & 4v + 2v_1 D_x^{-1} \\ 4v + 2v_1 D_x^{-1} & D_x^2 + 4u + 2u_1 D_x^{-1} \end{pmatrix} \\
\text{Root} & \begin{pmatrix} u_1 \\ v_1 \end{pmatrix}, \begin{pmatrix} v_1 \\ u_1 \end{pmatrix} \\
\text{Scaling} & \begin{pmatrix} xu_1 + 2u \\ xv_1 + 2v \end{pmatrix}
\end{array}$$

2.36 Coupled nonlinear wave system (Ito system)

Reference: [Ito82, AF87], [Dor93, p. 94];

$$\begin{array}{ll}
\text{Equation} & \begin{cases} u_t = u_3 + 6uu_1 + 2vv_1 \\ v_t = 2uv_1 + 2u_1v \end{cases} \\
\text{Hamiltonian} & \frac{u^2 + v^2}{2} \\
\text{Cosymplectic} & \begin{pmatrix} D_x^3 + 4uD_x + 2u_1 & 2vD_x \\ 2vD_x + 2v_1 & 0 \end{pmatrix} \\
\text{Symplectic} & \begin{pmatrix} D_x^{-1} & 0 \\ 0 & D_x^{-1} \end{pmatrix} \\
\text{Recursion} & \begin{pmatrix} D_x^2 + 4u + 2u_1 D_x^{-1} & 2v \\ 2v + 2v_1 D_x^{-1} & 0 \end{pmatrix} \\
\text{Root} & \begin{pmatrix} u_1 \\ v_1 \end{pmatrix} \\
\text{Scaling} & \begin{pmatrix} xu_1 + 2u \\ xv_1 + 2v \end{pmatrix}
\end{array}$$

2.37 Drinfel'd–Sokolov system

Reference: [Gök98];

$$\begin{array}{ll}
\text{Equation} & \begin{cases} u_t = 3vv_1 \\ v_t = 2v_3 + u_1v + 2uv_1 \end{cases} \\
\text{Hamiltonian} & \frac{v^2}{2} \\
\text{Cosymplectic} & \begin{pmatrix} 2D_x^3 + 2uD_x + u_1 & 2vD_x + v_1 \\ 2vD_x + v_1 & 2D_x^3 + 2uD_x + u_1 \end{pmatrix} \\
\text{Symplectic} & \begin{pmatrix} D_x^3 + \frac{5}{2}(uD_x + D_xu) & -15vD_x + \frac{15}{2}v_1 \\ -15vD_x - \frac{45}{2}v_1 & -27D_x^3 - \frac{27}{2}(uD_x + D_xu) \end{pmatrix} \\
& -\left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}^\dagger D_x^{-1} \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix}^\dagger + \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix}^\dagger D_x^{-1} \begin{pmatrix} 1 \\ 0 \end{pmatrix}^\dagger\right) \\
& -18 \begin{pmatrix} 0 \\ v \end{pmatrix}^\dagger D_x^{-1} \begin{pmatrix} 0 \\ v \end{pmatrix}^\dagger, \\
\text{Root} & \begin{pmatrix} u_1 \\ v_1 \end{pmatrix}; \begin{pmatrix} u_t \\ v_t \end{pmatrix}; \begin{pmatrix} h_1 \\ h_2 \end{pmatrix} \\
\text{Scaling} & \begin{pmatrix} xu_1 + 2u \\ xv_1 + 2v \end{pmatrix}
\end{array}$$

$$\eta_1 = -u_2 - 2u^2 + 3v^2,$$

$$\eta_2 = 9v_2 + 6uv,$$

$$h_1 = -2u_5 - 10uu_3 - 25u_1u_2 + 30vv_3 + 45v_1v_2 - 10u^2u_1 + 15v^2u_1 + 30uvv_1,$$

$$h_2 = 18v_5 + 10vu_3 + 35u_2v_1 + 45u_1v_2 + 30uv_3 + 10uu_1v + 10u^2v_1 + 15v^2v_1.$$

2.38 Benney system

Reference: [Ben73, AF87];

$$\begin{array}{ll} \text{Equation} & \begin{cases} u_t = vv_1 + 2D_x(uw) \\ v_t = 2u_1 + D_x(vw) \\ w_t = 2v_1 + 2ww_1 \end{cases} \\ \text{Hamiltonian} & uw + \frac{v^2}{2} \\ \text{Cosymplectic} & \begin{pmatrix} uD_x + D_xu & vD_x & wD_x \\ D_xv & 0 & 2D_x \\ D_xw & 2D_x & 0 \end{pmatrix} \\ \text{Symplectic} & \begin{pmatrix} 0 & 0 & D_x^{-1} \\ 0 & D_x^{-1} & 0 \\ D_x^{-1} & 0 & 0 \end{pmatrix} \\ \text{Recursion} & \begin{pmatrix} w & v & 2u + u_1D_x^{-1} \\ 2 & 0 & v + v_1D_x^{-1} \\ 0 & 2 & w + w_1D_x^{-1} \end{pmatrix} \\ \text{Root} & \begin{pmatrix} u_1 \\ v_1 \\ w_1 \end{pmatrix} \\ \text{Scaling} & \begin{pmatrix} xu_1 \\ xv_1 \\ xw_1 \end{pmatrix} + \alpha \begin{pmatrix} 3u \\ 2v \\ w \end{pmatrix}; \alpha \in \mathbb{C} \end{array}$$

2.39 Dispersive water wave system

Reference: [AF87];

$$\begin{array}{ll} \text{Equation} & \begin{cases} u_t = D_x(uw) \\ v_t = -v_2 + 2D_x(vw) + uu_1 \\ w_t = w_2 - 2v_1 + 2ww_1 \end{cases} \\ \text{Hamiltonian} & vw + \frac{u^2}{2} \\ \text{Cosymplectic} & \begin{pmatrix} 0 & D_xu & 0 \\ uD_x & vD_x + D_xv & -D_x^2 + wD_x \\ 0 & D_x^2 + D_xw & -2D_x \end{pmatrix} \\ \text{Symplectic} & \begin{pmatrix} D_x^{-1} & 0 & 0 \\ 0 & 0 & D_x^{-1} \\ 0 & D_x^{-1} & 0 \end{pmatrix} \\ \text{Recursion} & \begin{pmatrix} 0 & 0 & u + u_1D_x^{-1} \\ u & -D_x + w & 2v + v_1D_x^{-1} \\ 0 & -2 & D_x + w + w_1D_x^{-1} \end{pmatrix} \\ \text{Root} & \begin{pmatrix} u_1 \\ v_1 \\ w_1 \end{pmatrix} \end{array}$$

$$\text{Scaling} \quad \begin{pmatrix} xu_1 + \frac{3}{2}u \\ xv_1 + 2v \\ xw_1 + w \end{pmatrix}$$

If $u = 0$, this system reduces to the Broer–Kaup system studied in [Gök98].

References

- [AC91] Ablowitz M J and Clarkson P A, *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, Cambridge University Press, Cambridge, 1991.
- [AF87] Antonowicz M and Fordy A P, Coupled KdV equations with multi-Hamiltonian structures, *Phys. D* **28** (1987), 345–357.
- [Ben73] Benny D J, Some properties of long nonlinear waves, *Stud. Appl. Math.* **52** (1973), 45–50.
- [Bil93] Bilge A H, On the equivalence of linearization and formal symmetries as integrability tests for evolution equations, *J. Phys. A* **26**, No. 24 (1993), 7511–7519.
- [BPT83] Boiti M, Pempinelli F, and Tu G Z, The nonlinear evolution equations related to the Wadati–Konno–Ichikawa spectral problem, *Progr. Theoret. Phys.* **69**, No. 1 (1983), 48–64.
- [Cal87] Calogero F, The evolution partial differential equation $u_t = u_{xxx} + 3(u_{xx}u^2 + 3u_x^2u) + 3u_xu^4$, *J. Math. Phys.* **28**, No. 3 (1987), 538–555.
- [CD81] Calogero F and Degasperis A, Reduction technique for matrix nonlinear evolution equations solvable by the spectral transform, *J. Math. Phys.* **22** (1981), 23–31.
- [CDG76] Caudrey P J, Dodd R K and Gibbon J D, A new hierarchy of Korteweg–de Vries equation, *Proc. Roy. Soc. London Ser. A* **351** (1976), 407–422.
- [Cho87a] Chou T, Symmetries and a hierarchy of the general KdV equation, *J. Phys. A* **20** (1987), 359–366.
- [Cho87b] Chou T, Symmetries and a hierarchy of the general modified KdV equation, *J. Phys. A* **20** (1987), 367–374.
- [CLL87] Chen H H, Lee Y C and Lin J E, On the direct construction of the inverse scattering operators of integrable nonlinear Hamiltonian systems, *Phys. D* **26** (1987), 165–170.
- [CT88] Cavalcante J A and Tenenblat K, Conservation laws for nonlinear evolution equations, *J. Math. Phys.* **29**, No. 4 (1988), 1044–1049.
- [Dor93] Dorfman I, *Dirac structures and integrability of nonlinear evolution equations*, John Wiley & Sons Ltd., Chichester, 1993.
- [FF80] Fokas A S and Fuchssteiner B, On the structure of symplectic operators and hereditary symmetries, *Lett. al Nuovo Cimento* **28**, No. 8 (1980), 299–303.
- [FF81] Fuchssteiner B and Fokas A S, Symplectic structures, their Bäcklund transformations and hereditary symmetries, *Phys. D* **4** (1981), 47–60.
- [FG80] Fordy A P and Gibbons J, Integrable nonlinear Klein–Gordon equations and Toda Lattices, *Comm. Math. Phys.* **77** (1980), 21–30.

- [FG81] Fordy A P and Gibbons J, Factorization of operators. II, *J. Math. Phys.* **22**, No. 6 (1981), 1170–1175.
- [FO82] Fuchssteiner B and Oevel W, The bi-Hamiltonian structure of some nonlinear fifth- and seventh-order differential equations and recursion formulas for their symmetries and conserved covariants, *J. Math. Phys.* **23**, No. 3 (1982), 358–363.
- [Fok80] Fokas A S, A symmetry approach to exactly solvable evolution equations, *J. Math. Phys.* **21**, No. 6 (1980), 1318–1325.
- [FOW87] Fuchssteiner B, Oevel W and Wiwianka W, Computer-algebra methods for investigation of hereditary operators of higher order soliton equations, *Comput. Phys. Comm.* **44** (1987), 47–55.
- [Fuc79] Fuchssteiner B, Application of hereditary symmetries to nonlinear evolution equations, *Nonlinear Anal.* **3**, No. 11 (1979), 849–862.
- [Fuc82] Fuchssteiner B, The Lie algebra structure of degenerate Hamiltonian and bi-Hamiltonian systems, *Progr. Theoret. Phys.* **68**, No. 4 (1982), 1082–1104.
- [Gök98] Göktaş Ü, *Algorithmic Computation of Symmetries, Invariants and Recursion Operators for Systems of Nonlinear Evolution and Differential-Difference Equations*, PhD thesis, Colorado School of Mines, Golden, CO, 1998.
- [HS81] Hirota R and Satsuma J, Soliton solutions of a coupled Korteweg–de Vries equation, *Phys. Lett. A* **85** (1981), 407–408.
- [Ibr96] Ibragimov N H, editor, *CRC Handbook of Lie Group Analysis of Differential Equations*, volume 1-3. CRC Press, Inc, Boca Raton, 1996.
- [IŠ80] Ibragimov N H and Šabat A B, Infinite Lie-Bäcklund algebras, *Funktsional. Anal. i Prilozhen.* **14**, No. 4 (1980), 79–80.
- [Ito82] Ito M, Symmetries and conservation laws of a coupled nonlinear wave equation, *Phys. Lett. A* **91** (1982), 335–338.
- [Kap82] Kaptsov O V, Classification of evolution equations by conservation laws, *Funct. Anal. Appl.* **16**, No. 1 (1982), 59–61.
- [Kau80] Kaup D J, On the inverse scattering problem for cubic eigenvalue problems of the class $\psi_{xxx} + 6q\psi_x + 6r\psi = \lambda\psi$, *Stud. Appl. Math.* **62** (1980), 189–216.
- [KdV95] Korteweg D J and de Vries G, On the change of form of long waves advancing in a rectangular canal, and a new type of long stationary waves, *Philos. Mag.* **39**, No. 5 (1895), 422–443.
- [Kon87] Konopelchenko B G, *Nonlinear Integrable Equations*, Springer – New York, 1987.
- [Lax68] Lax P D, Integrals of nonlinear equations of evolution and solitary waves, *Comm. Pure Appl. Math.* **21** (1968), 467–490.
- [LLS⁺83] Leo M, Leo R A, Soliani G, Solombrino L and Martina L, Lie-Bäcklund symmetries for the Harry Dym equation, *Phys. Rev. D* **27**, No. 6 (1983), 1406–1408.
- [Ma93] Ma W X, A hierarchy of coupled Burgers systems possessing a hereditary structure, *J. Phys. A* **26**, No. 22 (1993), L1169–L1174.

- [Mag78] Magri F, A simple model of integrable Hamiltonian equation, *J. Math. Phys.* **19**, No. 5 (1978), 1156–1162.
- [MSS91] Mikhaïlov A V, Shabat A B and Sokolov V V, The symmetry approach to classification of integrable equations, In *What is integrability?*, Editor Zakharov V E, Springer – Berlin, 1991, 115–184.
- [Oev84] Oevel W, Rekursionsmechanismen für Symmetrien und Erhaltungssätze in integrablen Systemen, PhD thesis, Universität–Gesamthochschule Paderborn, Paderborn, 1984.
- [Oev90] Oevel G, Reduktion integrierbarer Systeme auf ihre Multisoliton Mannigfaltigkeiten, PhD thesis, Universität–Gesamthochschule Paderborn, Paderborn, 1990.
- [OF84] Oevel W and Fokas A S, Infinitely many commuting symmetries and constants of motion in involution for explicitly time-dependent evolution equations, *J. Math. Phys.* **25**, No. 4 (1984), 918–922.
- [Olv77] Olver P J, Evolution equations possessing infinitely symmetries, *J. Math. Phys.* **18**, No. 6 (1977), 1212–1215.
- [Olv93] Olver P J, Applications of Lie Groups to Differential Equations, second edition, Springer – New York, 1993.
- [SK74] Sawada K and Kotera T, A method of finding N-soliton solutions of the KdV and KdV-like equation, *Progr. Theoret. Phys.* **51** (1974), 1355–1367.
- [SW01a] Sanders J A and Wang J P, Integrable systems and their recursion operators, *Nonlinear Anal.* **47**, No. 8 (2001), 5213–5240.
- [SW01b] Sanders J A and Wang J P, On recursion operators, *Phys. D* **149**, No. 1-2 (2001), 1–10.
- [vBK91] van Bemmelen T and Kersten P, Nonlocal symmetries and recursion operators of the Landau–Lifshitz equation, *J. Math. Phys.* **32**, No. 7 (1991), 1709–1716.
- [Wan98] Wang J P, Symmetries and Conservation Laws of Evolution Equations, PhD thesis, Vrije Universiteit, Amsterdam, 1998.
- [WHV95] Willox R, Hereman W and Verheest F, Complete integrability of a modified vector derivative nonlinear Schrödinger equation, *Phys. Scripta* **52** (1995), 21–26.
- [WKI79] Wadati M, Konno K and Ichikawa Y H, New integrable nonlinear evolution equations, *J. Phys. Soc. Japan* **47**, No. 5 (1979), 1698–1700.
- [ZC86] Zhu G C and Chen H H, Symmetries and integrability of the cylindrical Korteweg–de Vries equation, *J. Math. Phys.* **27**, No. 1 (1986), 100–103.
- [ZS79] Zhiber A V and Shabat A B, Klein-Gordon equations with a nontrivial group, *Soviet Phys. Dokl.* **24** (1979), 607–609.
- [ZS01] Zhiber A V and Sokolov V V, Exactly integrable hyperbolic equations of Liouville type, *Uspehi mat. nauk* **56**, No. 1 (2001), 63–106.